arXiv:1008.2264v3 [math.FA] 26 Aug 2010

POINTWISE APPROXIMATION THEOREMS FOR COMBINATIONS OF BERNSTEIN POLYNOMIALS WITH INNER SINGULARITIES

WEN-MING LU AND LIN ZHANG

Abstract. We give direct and inverse theorems for the weighted approximation of functions with inner singularities by combinations of Bernstein polynomials.

1. Introduction

The set of all continuous functions, defined on the interval I, is denoted by $C(I)$. For any $f \in C([0,1])$, the corresponding *Bernstein operators* are defined as follows:

$$B_n(f,x) := \sum_{k=0}^{n} f(\frac{k}{n}) p_{n,k}(x),$$

where

$$p_{n,k}(x) := \binom{n}{k} x^k (1-x)^{n-k},\ k = 0,1,2,\dots,n,\ x \in [0,1].$$

Approximation properties of Bernstein operators have been studied very well (see [2], [4], [5]-[9], [14]-[16], for example). In order to approximate the functions with singularities, Della Vecchia et al. [4] and Yu-Zhao [14] introduced some kinds of modified Bernstein operators. Throughout the paper, C denotes a positive constant independent of n and x, which may be different in different cases. Ditzian and Totik [5] extended the method of combinations and defined the following combinations of Bernstein operators:

$$B_{n,r}(f,x) := \sum_{i=0}^{r-1} C_i(n) B_{n_i}(f,x),$$

with the conditions:

(a) $n = n_0 < n_1 < \cdots < n_{r-1} \leqslant Cn$,

(b) $\sum_{i=0}^{r-1} |C_i(n)| \leqslant C$,

(c) $\sum_{i=0}^{r-1} C_i(n) = 1$,

(d) $\sum_{i=0}^{r-1} C_i(n) n_i^{-k} = 0$, for $k = 1,\dots,r-1$.

For any positive integer r, we consider the determinant

$$A_r := \begin{vmatrix} 1 & 1 & 1 & \cdots & 1 \\ 2r+1 & 2r+2 & 2r+3 & \cdots & 4r+1 \\ (2r)(2r+1) & (2r+1)(2r+2) & (2r+2)(2r+3) & \cdots & (4r)(4r+1) \\ \cdots & \cdots & \cdots & \ddots & \cdots \\ 2\cdots(2r+1) & 3\cdots(2r+2) & 4\cdots(2r+3) & \cdots & (2r+2)\cdots(4r+1) \end{vmatrix}.$$

1991 *Mathematics Subject Classification.* Primary 41A10, Secondary 41A17.

Key words and phrases. Bernstein polynomials; Inner singularities; Pointwise approximation; Direct and inverse theorems.

We obtain $A_r = \prod_{j=2}^{2r} j!$. Thus, there is a unique solution for the system of nonhomogeneous linear equations:

$$\left\{\begin{array}{ccccccccc} a_1 & + & a_2 & + & \cdots & + & a_{2r+1} & = & 1, \\ (2r+1)a_1 & + & (2r+2)a_2 & + & \cdots & + & (4r+1)a_{2r+1} & = & 0, \\ (2r+1)(2r)a_1 & + & (2r+1)(2r+2)a_2 & + & \cdots & + & (4r)(4r+1)a_{2r+1} & = & 0, \\ & & & & \vdots & & & & \\ (2r+1)!a_1 & + & 3\cdots(2r+2)a_2 & + & \cdots & + & (2r+2)\cdots(4r+1)a_{2r+1} & = & 0. \end{array}\right. \tag{1.1}$$

Let

$$\psi(x) = \begin{cases} a_1x^{2r+1} + a_2x^{2r+2} + \cdots + a_{2r+1}x^{4r+1}, & 0 < x < 1, \\ 0, & x \leqslant 0, \\ 1, & x = 1. \end{cases}$$

with the coefficients $a_1, a_2, \cdots, a_{2r+1}$ satisfying (1.1). From (1.1), we see that $\psi(x) \in C^{(2r)}(-\infty, +\infty)$, $0 \leqslant \psi(x) \leqslant 1$ for $0 \leqslant x \leqslant 1$. Moreover, it holds that $\psi(1) = 1,\ \psi^{(i)}(0) = 0,\ i = 0, 1, \cdots, 2r$ and $\psi^{(i)}(1) = 0,\ i = 1, 2, \cdots, 2r$.

Let

$$H(f, x) := \sum_{i=1}^{r+1} f(x_i)l_i(x),$$

and

$$l_i(x) := \frac{\prod_{j=1,j\neq i}^{r+1}(x - x_j)}{\prod_{j=1,j\neq i}^{r+1}(x_i - x_j)},\ x_i = \frac{[n\xi - ((r-1)/2 + i)]}{n},\ i = 1, 2, \cdots r+1.$$

Further, let

$$x_1' = \frac{[n\xi - 2\sqrt{n}]}{n},\ x_2' = \frac{[n\xi - \sqrt{n}]}{n},\ x_3' = \frac{[n\xi + \sqrt{n}]}{n},\ x_4' = \frac{[n\xi + 2\sqrt{n}]}{n},$$

and

$$\bar{\psi}_1(x) = \psi(\frac{x - x_1'}{x_2' - x_1'}),\ \bar{\psi}_2(x) = \psi(\frac{x - x_3'}{x_4' - x_3'}).$$

Set

$$\bar{F}_n(f, x) := \bar{F}_n(x) = f(x)(1 - \bar{\psi}_1(x) + \bar{\psi}_2(x)) + \bar{\psi}_1(x)(1 - \bar{\psi}_2(x))H(x).$$

We have

$$\bar{F}_n(f, x) = \begin{cases} f(x), & x \in [0, x_{r-5/2}] \cup [x_{r+3/2}, 1], \\ f(x)(1 - \bar{\psi}_1(x)) + \bar{\psi}_1(x)H(x), & x \in [x_{r-5/2}, x_{r-3/2}], \\ H(x), & x \in [x_{r-3/2}, x_{r+1/2}], \\ H(x)(1 - \bar{\psi}_2(x)) + \bar{\psi}_2(x)f(x), & x \in [x_{r+1/2}, x_{r+3/2}]. \end{cases}$$

Obviously, $\bar{F}_n(f, x)$ is linear, reproduces polynomials of degree r, and $\bar{F}_n(f, x) \in C^{(2r)}([0, 1])$, provided that $f \in C^{(2r)}([0, 1])$. Now, we can define our new combinations of Bernstein operators as follows:

$$\bar{B}_{n,r}(f, x) := B_{n,r}(\bar{F}_n, x) = \sum_{i=0}^{r-1} C_i(n)B_{n_i}(\bar{F}_n, x), \tag{1.2}$$

where $C_i(n)$ satisfy the conditions (a)-(d).

2. THE MAIN RESULTS

Let $\phi : [0, 1] \longrightarrow R,\ \phi \neq 0$ be an admissible step-weight function of the Ditzian-Totik modulus of smoothness, that is, ϕ satisfies the following conditions:

(I) For every proper subinterval $[a, b] \subseteq [0, 1]$ there exists a constant $C_1 \equiv C(a, b) > 0$ such that $C_1^{-1} \leqslant \phi(x) \leqslant C_1$ for $x \in [a, b]$.

(II) There are two numbers $\beta(0) \geqslant 0$ and $\beta(1) \geqslant 0$ for which

$$\phi(x) \sim \begin{cases} x^{\beta(0)}, & \text{as } x \to 0+, \\ (1-x)^{\beta(1)}, & \text{as } x \to 1-. \end{cases}$$

($X \sim Y$ means $C^{-1}Y \leqslant X \leqslant CY$ for some C).

Combining conditions (**I**) and (**II**) on ϕ, we can deduce that

$$C^{-1}\phi_2(x) \leqslant \phi(x) \leqslant C\phi_2(x),\ x \in [0,1],$$

where $\phi_2(x) = x^{\beta(0)}(1-x)^{\beta(1)}$.

Let $\bar{w}(x) = |x-\xi|^{\alpha},\ 0<\xi<1,\ \alpha>0$ and $C_{\bar{w}} := \{f \in C([0,1]\setminus\xi) : \lim_{x\longrightarrow\xi}(\bar{w}f)(x) = 0\}$. The norm in $C_{\bar{w}}$ is defined as $\|f\|_{C_{\bar{w}}} := \|\bar{w}f\| = \sup_{0\leqslant x\leqslant 1}|(\bar{w}f)(x)|$. Define

$$W_{\phi}^{r} := \{f \in C_{\bar{w}} : f^{(r-1)} \in A.C.((0,1)),\ \|\bar{w}\phi^{r}f^{(r)}\| < \infty\},$$
$$W_{\varphi,\lambda}^{r} := \{f \in C_{\bar{w}} : f^{(r-1)} \in A.C.((0,1)),\ \|\bar{w}\varphi^{r\lambda}f^{(r)}\| < \infty\}.$$

For $f \in C_{\bar{w}}$, we define the *weighted modulus of smoothness* by

$$\omega_{\phi}^{r}(f,t)_{\bar{w}} := \sup_{0<h\leqslant t}\sup_{0\leqslant x\leqslant 1}|\bar{w}(x)\triangle_{h\phi(x)}^{r}f(x)|,$$

where

$$\Delta_{h\phi}^{r}f(x) = \sum_{k=0}^{r}(-1)^{k}\binom{r}{k}f(x+(\frac{r}{2}-k)h\phi(x)),$$
$$\overrightarrow{\Delta}_{h}^{r}f(x) = \sum_{k=0}^{r}(-1)^{k}\binom{r}{k}f(x+(r-k)h).$$

Recently Felten showed the following two theorems in [6]:

Theorem A. Let $\varphi(x) = \sqrt{x(1-x)}$ and let $\phi : [0,1] \longrightarrow R,\ \phi \neq 0$ be an admissible step-weight function of the Ditzian-Totik modulus of smoothness([5]) such that ϕ^2 and φ^2/ϕ^2 are concave. Then, for $f \in C[0,1]$ and $0<\alpha<2,\ |B_n(f,x)-f(x)| \leqslant \omega_{\phi}^{2}(f, n^{-1/2}\frac{\varphi(x)}{\phi(x)})$.

Theorem B. Let $\varphi(x) = \sqrt{x(1-x)}$ and let $\phi : [0,1] \longrightarrow R,\ \phi \neq 0$ be an admissible step-weight function of the Ditzian-Totik modulus of smoothness such that ϕ^2 and φ^2/ϕ^2 are concave. Then, for $f \in C[0,1]$ and $0<\alpha<2,\ |B_n(f,x)-f(x)| = O((n^{-1/2}\frac{\varphi(x)}{\phi(x)})^{\alpha})$ implies $\omega_{\phi}^{2}(f,t) = O(t^{\alpha})$.

Our main results are the following:

Theorem 2.1. *For any* $\alpha > 0,\ \min\{\beta(0),\beta(1)\} \geqslant \frac{1}{2},\ f \in C_{\bar{w}}$, *we have*

$$|\bar{w}(x)\phi^{r}(x)\bar{B}_{n,r-1}^{(r)}(f,x)| \leqslant Cn^{\frac{r}{2}}\|\bar{w}f\|. \tag{2.1}$$

Theorem 2.2. *For any* $\alpha > 0,\ f \in W_{\phi}^{r}$, *we have*

$$|\bar{w}(x)\phi^{r}(x)\bar{B}_{n,r-1}^{(r)}(f,x)| \leqslant C\|\bar{w}\phi^{r}f^{(r)}\|. \tag{2.2}$$

Theorem 2.3. *For* $f \in C_{\bar{w}},\ \alpha > 0,\ \min\{\beta(0),\beta(1)\} \geqslant \frac{1}{2},\ \alpha_0 \in (0,r)$, *we have*

$$\bar{w}(x)|f(x)-\bar{B}_{n,r-1}(f,x)| = O((n^{-\frac{1}{2}}\phi^{-1}(x)\delta_n(x))^{\alpha_0}) \Longleftrightarrow \omega_{\phi}^{r}(f,t)_{\bar{w}} = O(t^{\alpha_0}). \tag{2.3}$$

3. Lemmas

Lemma 3.1. *([15]) For any non-negative real u and v, we have*

$$\sum_{k=1}^{n-1}(\frac{k}{n})^{-u}(1-\frac{k}{n})^{-v}p_{n,k}(x) \leqslant Cx^{-u}(1-x)^{-v}. \tag{3.1}$$

Lemma 3.2. *([4]) If $\gamma \in R$, then*

$$\sum_{k=0}^{n} |k-nx|^{\gamma} p_{n,k}(x) \leqslant C n^{\frac{\gamma}{2}} \varphi^{\gamma}(x). \tag{3.2}$$

Lemma 3.3. *For any $f \in W_{\phi}^{r}$, $\alpha > 0$, we have*

$$\|\bar{w}\phi^{r} \bar{F}_{n}^{(r)}\| \leqslant C\|\bar{w}\phi^{r} f^{(r)}\|. \tag{3.3}$$

Proof. We first prove $x \in [x_{r-5/2}, x_{r-3/2}]$ (The same as the others), we have

$$\begin{aligned} |\bar{w}(x)\phi^{r}(x)\bar{F}_{n}^{(r)}(x)| &\leqslant |\bar{w}(x)\phi^{r}(x)f^{(r)}(x)| + |\bar{w}(x)\phi^{r}(x)(f(x)-\bar{F}_{n}(x))^{(r)}| \\ &:= I_1 + I_2. \end{aligned}$$

Obviously

$$I_1 \leqslant C\|\bar{w}\phi^{r} f^{(r)}\|.$$

For I_2, we have

$$I_2 = \bar{w}(x)\phi^{r}(x)|(f(x)-\bar{F}_{n}(x))^{(r)}| = \bar{w}(x)\phi^{r}(x)\sum_{i=0}^{r} n^{\frac{i}{2}}|(f(x)-\bar{F}_{n}(x))^{(r-i)}|.$$

By [5], we have

$$|(f(x)-\bar{F}_{n}(x))^{(r-i)}|_{[x_{r-5/2},x_{r-3/2}]} \leqslant C(n^{\frac{r-i}{2}}\|f-H\|_{[x_{r-5/2},x_{r-3/2}]} + n^{-\frac{i}{2}}\|f^{(r)}\|_{[x_{r-5/2},x_{r-3/2}]}).$$

So

$$\begin{aligned} I_2 &\leqslant C n^{\frac{r}{2}}\bar{w}(x)\phi^{r}(x)\|f-H\|_{[x_{r-5/2},x_{r-3/2}]} + C\bar{w}(x)\phi^{r}(x)\|f^{(r)}\|_{[x_{r-5/2},x_{r-3/2}]} \\ &:= T_1 + T_2. \end{aligned}$$

By Taylor expansion, we have

$$f(x_i) = \sum_{u=0}^{r-1} \frac{(x_i - x)^u}{u!} f^{(u)}(x) + \frac{1}{(r-1)!}\int_{x}^{x_i} (x_i - s)^{r-1} f^{(r)}(s)ds, \tag{3.4}$$

It follows from (3.4) and the identity

$$\sum_{i=1}^{r} x_i^{v} l_i(x) = C x^{v},\ v = 0, 1, \cdots, r.$$

we have

$$\begin{aligned} H(f,x) &= \sum_{i=1}^{r}\sum_{u=0}^{r} \frac{(x_i - x)^u}{u!} f^{(u)}(x) l_i(x) + \frac{1}{(r-1)!}\sum_{i=1}^{r} l_i(x)\int_{x}^{x_i}(x_i - s)^{r-1} f^{(r)}(s)ds \\ &= f(x) + C\sum_{u=1}^{r} f^{(u)}(x)(\sum_{v=0}^{u} C_{u}^{v}(-x)^{u-v}\sum_{i=1}^{r} x_i^{v} l_i(x)) \\ &\quad + \frac{1}{(r-1)!}\sum_{i=1}^{r} l_i(x)\int_{x}^{x_i}(x_i - s)^{r-1} f^{(r)}(s)ds, \end{aligned}$$

which implies that

$$\bar{w}(x)\phi^{r}(x)|f(x) - H(f,x)| = \frac{1}{(r-1)!}\bar{w}(x)\phi^{r}(x)\sum_{i=1}^{r} l_i(x)\int_{x}^{x_i}(x_i - s)^{r-1} f^{(r)}(s)ds,$$

since $|l_i(x)| \leqslant C$ for $x \in [x_{r-5/2}, x_{r-3/2}]$, $i = 1, 2, \cdots, r$. It follows from $\frac{|x_i-s|^{r-1}}{\bar{w}(s)} \leqslant \frac{|x_i-x|^{r-1}}{\bar{w}(x)}$, s between x_i and x, then

$$\begin{aligned}\bar{w}(x)\phi^r(x)|f(x)-H(f,x)| &= C\bar{w}(x)\phi^r(x)\sum_{i=1}^{r}\int_x^{x_i}(x_i-s)^{r-1}|f^{(r)}(s)|ds\\ &\leqslant C\phi^r(x)\|\bar{w}\phi^r f^{(r)}\|\sum_{i=1}^{r}(x_i-x)^{r-1}\int_x^{x_i}\phi^{-r}(s)ds\\ &\leqslant \frac{C}{n^{r/2}}\|\bar{w}\phi^r f^{(r)}\|.\end{aligned}$$

So

$$I_2 \leqslant C\|\bar{w}\phi^r f^{(r)}\|.$$

Then, the lemma is proved. □

Lemma 3.4. *If* $f \in W_\phi^r$, $\alpha > 0$, *then*

$$\bar{w}(x)|g(x)-H(g,x)| \leqslant C(\frac{\delta_n(x)}{\sqrt{n}\phi(x)})^r\|\bar{w}\phi^r g^{(r)}\|. \tag{3.5}$$

Proof. By Taylor expansion, we have

$$f(x_i) = \sum_{u=0}^{r-1}\frac{(x_i-x)^u}{u!}f^{(u)}(x) + \frac{1}{(r-1)!}\int_x^{x_i}(x_i-s)^{r-1}f^{(r)}(s)ds,$$

It follows from the above equality and the identity

$$\sum_{i=1}^{r}x_i^v l_i(x) = Cx^v,\ v = 0, 1, \cdots, r.$$

we have

$$\begin{aligned}H(f,x) &= \sum_{i=1}^{r}\sum_{u=0}^{r}\frac{(x_i-x)^u}{u!}f^{(u)}(x)l_i(x) + \frac{1}{(r-1)!}\sum_{i=1}^{r}l_i(x)\int_x^{x_i}(x_i-s)^{r-1}f^{(r)}(s)ds\\ &= f(x) + C\sum_{u=1}^{r}f^{(u)}(x)(\sum_{v=0}^{u}C_u^v(-x)^{u-v}\sum_{i=1}^{r}x_i^v l_i(x))\\ &\quad + \frac{1}{(r-1)!}\sum_{i=1}^{r}l_i(x)\int_x^{x_i}(x_i-s)^{r-1}f^{(r)}(s)ds,\end{aligned}$$

which implies that

$$\bar{w}(x)|g(x)-H(g,x)| = \frac{1}{(r-1)!}\bar{w}(x)\sum_{i=1}^{r}l_i(x)\int_x^{x_i}(x_i-s)^{r-1}g^{(r)}(s)ds,$$

since $|l_i(x)| \leqslant C$ for $x \in [x_{r-5/2}, x_{r-3/2}]$, $i = 1, 2, \cdots, r$. It follows from $\frac{|x_i-s|^{r-1}}{\bar{w}(s)} \leqslant \frac{|x_i-x|^{r-1}}{\bar{w}(x)}$, s between x_i and x, then

$$\begin{aligned}\bar{w}(x)|g(x)-H(g,x)| &\leqslant C\bar{w}(x)\sum_{i=1}^{r}\int_x^{x_i}(x_i-s)^{r-1}|g^{(r)}(s)|ds\\ &\leqslant C\|\bar{w}\phi^r g^{(r)}\|\sum_{i=1}^{r}(x_i-x)^{r-1}\int_x^{x_i}\phi^{-r}(s)ds\\ &\leqslant C\frac{\varphi^r(x)}{\phi^r(x)}\|\bar{w}\phi^r g^{(r)}\|\sum_{i=1}^{r}(x_i-x)^{r-1}\int_x^{x_i}\varphi^{-r}(s)ds\\ &\leqslant C\frac{\delta_n^r(x)}{\phi^r(x)}\|\bar{w}\phi^r g^{(r)}\|\sum_{i=1}^{r}(x_i-x)^{r-1}\int_x^{x_i}\varphi^{-r}(s)ds\\ &\leqslant C(\frac{\delta_n(x)}{\sqrt{n}\phi(x)})^r\|\bar{w}\phi^r g^{(r)}\|.\end{aligned}$$

$\square$

Lemma 3.5. *For any* $\alpha > 0$, $f \in C_{\bar{w}}$, *we have*

$$\|\bar{w}\bar{B}_{n,r-1}(f)\| \leqslant C\|\bar{w}f\|. \tag{3.6}$$

Proof. By (1.2), we have

$$\begin{aligned} |\bar{w}(x)\bar{B}_{n,r}(f,x)| &= |\bar{w}(x)B_{n,r}(\bar{F}_n,x)| \leqslant \bar{w}(x)\sum_{i=0}^{r-1}\sum_{k=1}^{n_i-1}C_i(n)|\bar{F}_n(\frac{k}{n_i})|p_{n_i,k}(x) \\ &\quad +\bar{w}(x)\sum_{i=0}^{r-1}C_i(n)|\bar{F}_n(0)|p_{n_i,0}(x)+\bar{w}(x)\sum_{i=0}^{r-1}C_i(n)|\bar{F}_n(1)|p_{n_i,n_i}(x) \\ &:= I_1+I_2+I_3. \end{aligned}$$

Now, the theorem can be proved easily. $\square$

Lemma 3.6. *([17]) Let* $\min\{\beta(0),\beta(1)\} \geqslant \frac{1}{2}$, *then for* $r \in N$, $0 < t < \frac{1}{8r}$ *and* $\frac{rt}{2} < x < 1-\frac{rt}{2}$, *we have*

$$\int_{-\frac{t}{2}}^{\frac{t}{2}}\cdots\int_{-\frac{t}{2}}^{\frac{t}{2}}\phi^{-r}(x+\sum_{k=1}^{r}u_k)du_1\cdots du_r \leqslant Ct^r\phi^{-r}(x). \tag{3.7}$$

Lemma 3.7. *Let* $A_n(x) := \bar{w}(x)\sum_{|k-n\xi|\leqslant\sqrt{n}}p_{n,k}(x)$. *Then* $A_n(x) \leqslant Cn^{-\alpha/2}$ *for* $0<\xi<1$ *and* $\alpha>0$.

Proof. If $|x-\xi| \leqslant \frac{3}{\sqrt{n}}$, then the statement is trivial. Hence assume $0 \leqslant x \leqslant \xi-\frac{3}{\sqrt{n}}$ (the case $\xi+\frac{3}{\sqrt{n}} \leqslant x \leqslant 1$ can be treated similarly). Then for a fixed x the maximum of $p_{n,k}(x)$ is attained for $k=k_n:=[n\xi-\sqrt{n}]$. By using Stirling's formula, we get

$$\begin{aligned} p_{n,k_n}(x) &\leqslant C\frac{(\frac{n}{e})^n\sqrt{n}x^{k_n}(1-x)^{n-k_n}}{(\frac{k_n}{e})^{k_n}\sqrt{k_n}(\frac{n-k_n}{e})^{n-k_n}\sqrt{n-k_n}} \\ &\leqslant \frac{C}{\sqrt{n}}(\frac{nx}{k_n})^{k_n}(\frac{n(1-x)}{n-k_n})^{n-k_n} \\ &= \frac{C}{\sqrt{n}}(1-\frac{k_n-nx}{k_n})^{k_n}(1+\frac{k_n-nx}{n-k_n})^{n-k_n}. \end{aligned}$$

Now from the inequalities

$$k_n-nx=[n\xi-\sqrt{n}]-nx > n(\xi-x)-\sqrt{n}-1 \geqslant \frac{1}{2}n(\xi-x),$$

and

$$1-u \leqslant e^{-u-\frac{1}{2}u^2},\ 1+u \leqslant e^u,\ u \geqslant 0.$$

We have that the second inequality is valid. To prove the first one we consider the function $\lambda(u)=e^{-u-\frac{1}{2}u^2}+u-1$. Here $\lambda(0)=0$, $\lambda'(u)=-(1+u)e^{-u-\frac{1}{2}u^2}+1$, $\lambda'(0)=0$, $\lambda''(u)=u(u+2)e^{-u-\frac{1}{2}u^2} \geqslant 0$, whence $\lambda(u) \geqslant 0$ for $u \geqslant 0$. Hence

$$\begin{aligned} p_{n,k_n}(x) &\leqslant \frac{C}{\sqrt{n}}exp\{k_n[-\frac{k_n-nx}{k_n}-\frac{1}{2}(\frac{k_n-nx}{k_n})^2]+k_n-nx\} \\ &= \frac{C}{\sqrt{n}}exp\{-\frac{(k_n-nx)^2}{2k_n}\} \leqslant e^{-Cn(\xi-x)^2}. \end{aligned}$$

Thus $A_n(x) \leqslant C(\xi-x)^\alpha e^{-Cn(\xi-x)^2}$. An easy calculation shows that here the maximum is attained when $\xi-x=\frac{C}{\sqrt{n}}$ and the lemma follows. $\square$

Lemma 3.8. *For* $0<\xi<1$, α, $\beta>0$, *we have*

$$\bar{w}(x)\sum_{|k-n\xi|\leqslant\sqrt{n}}|k-nx|^\beta p_{n,k}(x) \leqslant Cn^{\frac{\beta-\alpha}{2}}\varphi^\beta(x). \tag{3.8}$$

Proof. By (3.2) and the lemma 3.7, we have

$$\bar{w}(x)^{\frac{1}{2n}}(\bar{w}(x)\sum_{|k-n\xi|\leqslant\sqrt{n}}p_{n,k}(x))^{\frac{2n-1}{2n}}(\sum_{|k-n\xi|\leqslant\sqrt{n}}|k-nx|^{2n\beta}p_{n,k}(x))^{\frac{1}{2n}}\leqslant Cn^{\frac{\beta-\alpha}{2}}\varphi^{\beta}(x).$$

□

Lemma 3.9. *For any* $\alpha>0$, $f\in W_{\phi}^{r}$, *we have*

$$\|\bar{w}\bar{B}_{n,r-1}^{(r)}(f)\|\leqslant Cn^{r}\|\bar{w}f\|. \tag{3.9}$$

Proof. We first prove $x\in[0,\frac{1}{n})$ (The same as $x\in(1-\frac{1}{n},1]$), now

$$\begin{aligned}
|\bar{w}(x)\bar{B}_{n,r-1}^{(r)}(f,x)| &\leqslant \bar{w}(x)\sum_{i=0}^{r-2}\frac{n_i!}{(n_i-r)!}\sum_{k=0}^{n_i-r}C_i(n)|\overrightarrow{\Delta}_{\frac{1}{n_i}}^{r}\bar{F}_n(\frac{k}{n_i})|p_{n_i-r,k}(x)\\
&\leqslant C\bar{w}(x)\sum_{i=0}^{r-2}n_i^{r}\sum_{k=0}^{n_i-r}|\overrightarrow{\Delta}_{\frac{1}{n_i}}^{r}\bar{F}_n(\frac{k}{n_i})|p_{n_i-r,k}(x)\\
&\leqslant C\bar{w}(x)\sum_{i=0}^{r-2}n_i^{r}\sum_{k=0}^{n_i-r}\sum_{j=0}^{r}C_r^{j}|\bar{F}_n(\frac{k+r-j}{n_i})|p_{n_i-r,k}(x)\\
&\leqslant C\bar{w}(x)\sum_{i=0}^{r-2}n_i^{r}\sum_{j=0}^{r}C_r^{j}|\bar{F}_n(\frac{r-j}{n_i})|p_{n_i-r,0}(x)\\
&\quad + C\bar{w}(x)\sum_{i=0}^{r-2}n_i^{r}\sum_{j=0}^{r}C_r^{j}|\bar{F}_n(\frac{n_i-j}{n_i})|p_{n_i-r,n_i-r}(x)\\
&\quad + C\bar{w}(x)\sum_{i=0}^{r-2}n_i^{r}\sum_{k=1}^{n_i-r-1}\sum_{j=0}^{r}C_r^{j}|\bar{F}_n(\frac{k+r-j}{n_i})|p_{n_i-r,k}(x)\\
&:= H_1+H_2+H_3.
\end{aligned}$$

We have

$$\begin{aligned}
H_1 &\leqslant C\bar{w}(x)\sum_{i=0}^{r-2}n_i^{r}\sum_{j=0}^{r}|\bar{F}_n(\frac{r-j}{n_i})|p_{n_i-r,0}(x)\\
&\leqslant Cn^{r}\|\bar{w}f\|\sum_{i=0}^{r-2}\sum_{j=0}^{r}(\frac{n_i|x-\xi|}{r-j-n_i\xi})^{\alpha}(1-x)^{n_i-r}\\
&\leqslant Cn^{r}\|\bar{w}f\|\sum_{i=0}^{r-2}(n_i|x-\xi|)^{\alpha}(1-x)^{n_i-r}\\
&\leqslant Cn^{r}\|\bar{w}f\|.
\end{aligned}$$

Similarly, we can get $H_2\leqslant Cn^{r}\|\bar{w}f\|$, and $H_3\leqslant Cn^{r}\|\bar{w}f\|$.

When $x\in[\frac{1}{n},1-\frac{1}{n}]$, according to [5], we have

$$\begin{aligned}
&|\bar{w}(x)\bar{B}_{n,r-1}^{(r)}(f,x)|\\
=\ &|\bar{w}(x)B_{n,r-1}^{(r)}(\bar{F}_n,x)|\\
\leqslant\ &\bar{w}(x)(\varphi^{2}(x))^{-r}\sum_{i=0}^{r-2}\sum_{j=0}^{r}Q_j(x,n_i)C_i(n)n_i^{j}\sum_{k/n_i\in A}|(x-\frac{k}{n_i})^{j}||\bar{F}_n(\frac{k}{n_i})|p_{n_i,k}(x)\\
&+\bar{w}(x)(\varphi^{2}(x))^{-r}\sum_{i=0}^{r-2}\sum_{j=0}^{r}Q_j(x,n_i)C_i(n)n_i^{j}\sum_{x_2'\leqslant k/n_i\leqslant x_3'}|(x-\frac{k}{n_i})^{j}||H(\frac{k}{n_i})|p_{n_i,k}(x)\\
:=\ &\sigma_1+\sigma_2.
\end{aligned}$$

Where $A := [0, x_2'] \cup [x_3', 1]$, H is a linear function. If $\frac{k}{n}_i \in A$, when $\frac{\bar{w}(x)}{\bar{w}(\frac{k}{n_i})} \leqslant C(1 + n_i^{-\frac{\alpha}{2}}|k - n_i x|^{\alpha})$, we have $|k - n_i\xi| \geqslant \frac{\sqrt{n_i}}{2}$, also $Q_j(x, n_i) = (n_i x(1-x))^{[\frac{r-j}{2}]}$, and $(\varphi^2(x))^{-r} Q_j(x, n_i) n_i^j \leqslant C(n_i/\varphi^2(x))^{\frac{r+j}{2}}$. By (3.2), then

$$\begin{aligned}
\sigma_1 &\leqslant C\bar{w}(x)\sum_{i=0}^{r-2}\sum_{j=0}^{r}\left(\frac{n_i}{\varphi^2(x)}\right)^{\frac{r+j}{2}}\sum_{k=0}^{n_i}|(x - \frac{k}{n}_i)^j||\bar{F}_n(\frac{k}{n}_i)|p_{n_i,k}(x) \\
&\leqslant C\|\bar{w}f\|\sum_{i=0}^{r-2}\sum_{j=0}^{r}\left(\frac{n_i}{\varphi^2(x)}\right)^{\frac{r+j}{2}}\sum_{k=0}^{n_i}[1 + n_i^{-\frac{\alpha}{2}}|k - n_i x|^{\alpha}]|x - \frac{k}{n}_i|^j p_{n_i,k}(x) \\
&:= I_1 + I_2.
\end{aligned}$$

By a simple calculation, we have $I_1 \leqslant Cn^r\|\bar{w}f\|$. By (3.2), then

$$I_2 \leqslant C\|\bar{w}f\|\sum_{i=0}^{r-2}\sum_{j=0}^{r} n_i^{-(\frac{\alpha}{2}+j)}\left(\frac{n_i}{\varphi^2(x)}\right)^{\frac{r+j}{2}}\sum_{k=0}^{n_i}|k - n_i x|^{\alpha+j}p_{n_i,k}(x) \leqslant Cn^r\|\bar{w}f\|.$$

We note that $|H(\frac{k}{n_i})| \leqslant max(|H(x_1')|, |H(x_4')|) := H(a)$.

If $x \in [x_1', x_4']$, we have $\bar{w}(x) \leqslant \bar{w}(a)$. So, if $x \in [x_1', x_4']$, then

$$\sigma_2 \leqslant Cn^r\bar{w}(a)H(a) \leqslant Cn^r\|\bar{w}f\|.$$

If $x \notin [x_1', x_4']$, then $\bar{w}(a) > n_i^{-\frac{\alpha}{2}}$, by (3.8), we have

$$\begin{aligned}
\sigma_2 \leqslant C\bar{w}(a)H(a)\bar{w}(x)\sum_{i=0}^{r-2}\sum_{j=0}^{r} n_i^{\frac{\alpha}{2}}\left(\frac{n_i}{\varphi^2(x)}\right)^{\frac{r+j}{2}}\sum_{x_2' \leqslant k/n_i \leqslant x_3'}|x - \frac{k}{n}_i|^j p_{n_i,k}(x) \\
\leqslant Cn^r\|\bar{w}f\|.
\end{aligned}$$

It follows from combining the above inequalities that the lemma is proved. $\square$

4. PROOF OF THEOREMS

4.1. **Proof of Theorem 2.1.** When $f \in C_{\bar{w}},\ \min\{\beta(0), \beta(1)\} \geqslant \frac{1}{2}$, we discuss it as follows:

Case 1. If $0 \leqslant \varphi(x) \leqslant \frac{1}{\sqrt{n}}$, by (3.9), we have

$$\begin{aligned}
|\bar{w}(x)\phi^r(x)\bar{B}_{n,r-1}^{(r)}(f, x)| &= C\varphi^r(x)\cdot\frac{\phi^r(x)}{\varphi^r(x)}|\bar{w}(x)\bar{B}_{n,r-1}^{(r)}(f, x)| \\
&\leqslant Cn^{\frac{r}{2}}\|\bar{w}f\|. \qquad (4.1)
\end{aligned}$$

Case 2. Case 2. If $\varphi(x) > \frac{1}{\sqrt{n}}$, we have

$$\begin{aligned}
&|\bar{B}_{n,r-1}^{(r)}(f, x)| = |B_{n,r-1}^{(r)}(\bar{F}_n, x)| \\
\leqslant\ &(\varphi^2(x))^{-r}\sum_{i=0}^{r-2}\sum_{j=0}^{r} Q_j(x, n_i)C_i(n)n_i^j\sum_{k=0}^{n_i}|(x - \frac{k}{n}_i)^j||\bar{F}_n(\frac{k}{n}_i)|p_{n_i,k}(x),
\end{aligned}$$

where
$Q_j(x, n_i) = (n_i x(1-x))^{[\frac{r-j}{2}]}$, and $(\varphi^2(x))^{-r}Q_j(x, n_i)n_i^j \leqslant C(n_i/\varphi^2(x))^{\frac{r+j}{2}}$.

So

$$\begin{aligned}
&|\bar{w}(x)\phi^r(x)\bar{B}^{(r)}_{n,r-1}(f,x)|\\
\leqslant\ & C\bar{w}(x)\phi^r(x)\sum_{i=0}^{r-2}\sum_{j=0}^{r}(\frac{n_i}{\varphi^2(x)})^{\frac{r+j}{2}}\sum_{k=0}^{n_i}|(x-\frac{k}{n_i})^j||\bar{F}_n(\frac{k}{n_i})|p_{n_i,k}(x)\\
=\ & C\bar{w}(x)\phi^r(x)\sum_{i=0}^{r-2}\sum_{j=0}^{r}(\frac{n_i}{\varphi^2(x)})^{\frac{r+j}{2}}\sum_{k/n_i\in A}|(x-\frac{k}{n_i})^j||\bar{F}_n(\frac{k}{n_i})|p_{n_i,k}(x)\\
& +C\bar{w}(x)\phi^r(x)\sum_{i=0}^{r-2}\sum_{j=0}^{r}(\frac{n_i}{\varphi^2(x)})^{\frac{r+j}{2}}\sum_{x_2'\leqslant k/n_i\leqslant x_3'}|(x-\frac{k}{n_i})^j||H(\frac{k}{n_i})|p_{n_i,k}(x)\\
:=\ & \sigma_1+\sigma_2. \qquad (4.2)
\end{aligned}$$

Where $A:=[0,x_2']\cup[x_3',1]$, we can easily get $\sigma_1\leqslant Cn^{\frac{r}{2}}\|\bar{w}f\|$, and $\sigma_2\leqslant Cn^{\frac{r}{2}}\|\bar{w}f\|$. By bringing these facts together, the theorem is proved. □

4.2. **Proof of Theorem 2.2.** When $f\in W^r_{\phi}$, by [5], we have

$$B^{(r)}_{n,r-1}(\bar{F}_n,x)=\sum_{i=0}^{r-2}C_i(n)n_i^r\sum_{k=0}^{n_i-r}\overrightarrow{\Delta}^r_{\frac{1}{n_i}}\bar{F}_n(\frac{k}{n_i})p_{n_i-r,k}(x). \tag{4.3}$$

If $0<k<n_i-r$, we have

$$|\overrightarrow{\Delta}^r_{\frac{1}{n_i}}\bar{F}_n(\frac{k}{n_i})|\leqslant Cn_i^{-r+1}\int_0^{\frac{r}{n_i}}|\bar{F}_n^{(r)}(\frac{k}{n_i}+u)|du, \tag{4.4}$$

If $k=0$, we have

$$|\overrightarrow{\Delta}^r_{\frac{1}{n_i}}\bar{F}_n(0)|\leqslant C\int_0^{\frac{r}{n_i}}u^{r-1}|\bar{F}_n^{(r)}(u)|du, \tag{4.5}$$

Similarly

$$|\overrightarrow{\Delta}^r_{\frac{1}{n_i}}\bar{F}_n(\frac{n_i-r}{n_i})|\leqslant Cn_i^{-r+1}\int_{1-\frac{r}{n_i}}^{1}(1-u)^{\frac{r}{2}}|\bar{F}_n^{(r)}(u)|du. \tag{4.6}$$

By (4.3), we have

$$\begin{aligned}
&|\bar{w}(x)\phi^r(x)\bar{B}^{(r)}_{n,r-1}(f,x)|\\
&\leqslant C\bar{w}(x)\phi^r(x)\sum_{i=0}^{r-2}n_i^r\sum_{k=0}^{n_i-r}|\overrightarrow{\Delta}^r_{\frac{1}{n_i}}\bar{F}_n(\frac{k}{n_i})|p_{n_i-r,k}(x)\\
&\leqslant C\bar{w}(x)\phi^r(x)\sum_{i=0}^{r-2}n_i^r\sum_{k=1}^{n_i-r-1}|\overrightarrow{\Delta}^r_{\frac{1}{n_i}}\bar{F}_n(\frac{k}{n_i})|p_{n_i-r,k}(x)\\
&+C\bar{w}(x)\phi^r(x)\sum_{i=0}^{r-2}n_i^r|\overrightarrow{\Delta}^r_{\frac{1}{n_i}}\bar{F}_n(0)|p_{n_i-r,0}(x)\\
&+C\bar{w}(x)\phi^r(x)\sum_{i=0}^{r-2}n_i^r|\overrightarrow{\Delta}^r_{\frac{1}{n_i}}\bar{F}_n(1)|p_{n_i-r,n_i-r}(x). \qquad (4.7)
\end{aligned}$$

which combining with (4.4)-(4.6) give

$$|\bar{w}(x)\phi^r(x)\bar{B}^{(r)}_{n,r-1}(f,x)|\leqslant C\|w\phi^rf^{(r)}\|.\square$$

Combining with the theorem 2.1 and theorem 2.2, we can obtain

Corollary *For any* $\alpha>0,\ 0\leqslant\lambda\leqslant1,$ *we have*

$$|\bar{w}(x)\varphi^{r\lambda}(x)\bar{B}^{(r)}_{n,r-1}(f,x)|\leqslant\left\{\begin{array}{ll}Cn^{r/2}\{max\{n^{r(1-\lambda)/2},\varphi^{r(\lambda-1)}(x)\}\}\|\bar{w}f\|, & f\in C_{\bar{w}},\\ C\|\bar{w}\varphi^{r\lambda}f^{(r)}\|, & f\in W^r_{\bar{w},\lambda}.\end{array}\right. \tag{4.8}$$

4.3. **Proof of Theorem 2.3.**

4.3.1. *The direct theorem.* We know

$$\bar{F}_n(t) = \bar{F}_n(x) + \bar{F}'_n(t)(t-x) + \cdots + \frac{1}{(r-1)!}\int_x^t (t-u)^{r-1}\bar{F}_n^{(r)}(u)du, \tag{4.9}$$

$$B_{n,r-1}((\cdot - x)^k, x) = 0,\ k = 1, 2, \cdots, r-1. \tag{4.10}$$

According to the definition of W_ϕ^r, for any $g \in W_\phi^r$, we have $\bar{B}_{n,r-1}(g,x) = B_{n,r-1}(\bar{G}_n(g),x)$, and $\bar{w}(x)|\bar{G}_n(x) - B_{n,r-1}(\bar{G}_n,x)| = \bar{w}(x)|B_{n,r-1}(R_r(\bar{G}_n,t,x),x)|$, thereof $R_r(\bar{G}_n,t,x) = \int_x^t (t-u)^{r-1}\bar{G}_n^{(r)}(u)du$, we have

$$\begin{aligned}\bar{w}(x)|\bar{G}_n(x) - B_{n,r-1}(\bar{G}_n,x)| &\leqslant C\|\bar{w}\phi^r\bar{G}_n^{(r)}\|\bar{w}(x)B_{n,r-1}(\int_x^t \frac{|t-u|^{r-1}}{\bar{w}(u)\phi^r(u)}du, x)\\ &\leqslant C\|\bar{w}\phi^r\bar{G}_n^{(r)}\|\bar{w}(x)(B_{n,r-1}(\int_x^t \frac{|t-u|^{r-1}}{\phi^{2r}(u)}du, x))^{\frac{1}{2}}\cdot\\ &(B_{n,r-1}(\int_x^t \frac{|t-u|^{r-1}}{\bar{w}^2(u)}du, x))^{\frac{1}{2}}.\end{aligned} \tag{4.11}$$

also

$$\int_x^t \frac{|t-u|^{r-1}}{\phi^{2r}(u)}du \leqslant C\frac{|t-x|^r}{\phi^{2r}(x)},\ \int_x^t \frac{|t-u|^{r-1}}{\bar{w}^2(u)}du \leqslant C\frac{|t-x|^r}{\bar{w}^2(x)}. \tag{4.12}$$

By (3.2), (3.3) and (4.12), we have

$$\begin{aligned}\bar{w}(x)|\bar{G}_n(x) - B_{n,r-1}(\bar{G}_n,x)| &\leqslant C\|\bar{w}\phi^r\bar{G}_n^{(r)}\|\phi^{-r}(x)B_{n,r-1}(|t-x|^r, x)\\ &\leqslant Cn^{-\frac{r}{2}}\frac{\varphi^r(x)}{\phi^r(x)}\|\bar{w}\phi^r\bar{G}_n^{(r)}\|\\ &\leqslant Cn^{-\frac{r}{2}}\frac{\delta_n^r(x)}{\phi^r(x)}\|\bar{w}\phi^r\bar{G}_n^{(r)}\|\\ &= C(\frac{\delta_n(x)}{\sqrt{n}\phi(x)})^r\|\bar{w}\phi^r\bar{G}_n^{(r)}\|.\end{aligned} \tag{4.13}$$

By (3.3), (3.5) and (4.13), when $g \in W_\phi^r$, then

$$\begin{aligned}\bar{w}(x)|g(x) - \bar{B}_{n,r-1}(g,x)| &\leqslant \bar{w}(x)|g(x) - \bar{G}_n(g,x)| + \bar{w}(x)|\bar{G}_n(g,x) - \bar{B}_{n,r-1}(g,x)|\\ &\leqslant \bar{w}(x)|g(x) - H(g,x)|\\ &+\ C(\frac{\delta_n(x)}{\sqrt{n}\phi(x)})^r\|\bar{w}\phi^r\bar{G}_n^{(r)}\|\\ &\leqslant C(\frac{\delta_n(x)}{\sqrt{n}\phi(x)})^r\|\bar{w}\phi^r g^{(r)}\|.\end{aligned} \tag{4.14}$$

For $f \in C_{\bar{w}}$, we choose proper $g \in W_\phi^r$, by (3.6) and (4.14), then

$$\begin{aligned}\bar{w}(x)|f(x) - \bar{B}_{n,r-1}(f,x)| &\leqslant \bar{w}(x)|f(x) - g(x)| + \bar{w}(x)|\bar{B}_{n,r-1}(f-g,x)|\\ &+\bar{w}(x)|g(x) - \bar{B}_{n,r-1}(g,x)|\\ &\leqslant C(\|\bar{w}(f-g)\| + (\frac{\delta_n(x)}{\sqrt{n}\phi(x)})^r\|\bar{w}\phi^r g^{(r)}\|)\\ &\leqslant C\omega_\phi^r(f, \frac{\delta_n(x)}{\sqrt{n}\phi(x)})_{\bar{w}}.\Box\end{aligned}$$

4.3.2. *The inverse theorem.* The *weighted K-function* is given by

$$K_{r,\phi}(f,t^r)_{\bar{w}} := \inf_g \{\|\bar{w}(f-g)\| + t^r\|\bar{w}\phi^r g^{(r)}\| : g \in W_\phi^r\}.$$

By [5], we have

$$C^{-1}\omega_\phi^r(f,t)_{\bar{w}} \leqslant K_{r,\phi}(f,t^r)_{\bar{w}} \leqslant C\omega_\phi^r(f,t)_{\bar{w}}. \tag{4.15}$$

Proof. Let $\delta > 0$, by (4.15), we choose proper g so that

$$\|\bar{w}(f-g)\| \leqslant C\omega_\phi^r(f,\delta)_{\bar{w}},\ \|\bar{w}\phi^r g^{(r)}\| \leqslant C\delta^{-r}\omega_\phi^r(f,\delta)_{\bar{w}}. \tag{4.16}$$

For $r \in N,\ 0 < t < \frac{1}{8r}$ and $\frac{rt}{2} < x < 1 - \frac{rt}{2}$, we have

$$\begin{aligned}
|\bar{w}(x)\Delta_{h\phi}^r f(x)| &\leqslant |\bar{w}(x)\Delta_{h\phi}^r(f(x) - \bar{B}_{n,r-1}(f,x))| + |\bar{w}(x)\Delta_{h\phi}^r\bar{B}_{n,r-1}(f-g,x)| \\
&\quad + |\bar{w}(x)\Delta_{h\phi}^r\bar{B}_{n,r-1}(g,x)| \\
&\leqslant \sum_{j=0}^r C_r^j \Big(n^{-\frac{1}{2}}\frac{\delta_n(x+(\frac{r}{2}-j)h\phi(x))}{\phi(x+(\frac{r}{2}-j)h\phi(x))}\Big)^{\alpha_0} \\
&\quad + \int_{-\frac{h\phi(x)}{2}}^{\frac{h\phi(x)}{2}} \cdots \int_{-\frac{h\phi(x)}{2}}^{\frac{h\phi(x)}{2}} \bar{w}(x)\bar{B}_{n,r-1}^{(r)}(f-g, x+\sum_{k=1}^r u_k)du_1\cdots du_r \\
&\quad + \int_{-\frac{h\phi(x)}{2}}^{\frac{h\phi(x)}{2}} \cdots \int_{-\frac{h\phi(x)}{2}}^{\frac{h\phi(x)}{2}} \bar{w}(x)\bar{B}_{n,r-1}^{(r)}(g, x+\sum_{k=1}^r u_k)du_1\cdots du_r \\
&:= J_1 + J_2 + J_3.
\end{aligned} \tag{4.17}$$

Obviously

$$J_1 \leqslant C(n^{-\frac{1}{2}}\phi^{-1}(x)\delta_n(x))^{\alpha_0}. \tag{4.18}$$

By (3.9) and (4.16), we have

$$\begin{aligned}
J_2 &\leqslant Cn^r\|\bar{w}(f-g)\| \int_{-\frac{h\phi(x)}{2}}^{\frac{h\phi(x)}{2}} \cdots \int_{-\frac{h\phi(x)}{2}}^{\frac{h\phi(x)}{2}} du_1\cdots du_r \\
&\leqslant Cn^r h^r\phi^r(x)\|\bar{w}(f-g)\| \\
&\leqslant Cn^r h^r\phi^r(x)\omega_\phi^r(f,\delta)_{\bar{w}}.
\end{aligned} \tag{4.19}$$

By the first inequality of (4.8), we let $\lambda = 1$, and (4.16), then

$$\begin{aligned}
J_2 &\leqslant Cn^{\frac{r}{2}}\|\bar{w}(f-g)\| \int_{-\frac{h\phi(x)}{2}}^{\frac{h\phi(x)}{2}} \cdots \int_{-\frac{h\phi(x)}{2}}^{\frac{h\phi(x)}{2}} \varphi^{-r}(x+\sum_{k=1}^r u_k)du_1\cdots du_r \\
&\leqslant Cn^{\frac{r}{2}}h^r\phi^r(x)\varphi^{-r}(x)\|\bar{w}(f-g)\| \\
&\leqslant Cn^{\frac{r}{2}}h^r\phi^r(x)\varphi^{-r}(x)\omega_\phi^r(f,\delta)_{\bar{w}}.
\end{aligned} \tag{4.20}$$

By (3.7) and (4.16), we have

$$\begin{aligned}
J_3 &\leqslant C\|\bar{w}\phi^r g^{(r)}\|\bar{w}(x) \int_{-\frac{h\phi(x)}{2}}^{\frac{h\phi(x)}{2}} \cdots \int_{-\frac{h\phi(x)}{2}}^{\frac{h\phi(x)}{2}} \bar{w}^{-1}(x+\sum_{k=1}^r u_k)\phi^{-r}(x+\sum_{k=1}^r u_k)du_1\cdots du_r \\
&\leqslant Ch^r\|\bar{w}\phi^r g^{(r)}\| \\
&\leqslant Ch^r\delta^{-r}\omega_\phi^r(f,\delta)_{\bar{w}}.
\end{aligned} \tag{4.21}$$

Now, by (4.17)-(4.21), there exists a constant $M > 0$ so that

$$\begin{aligned} |\bar{w}(x)\Delta_{h\phi}^r f(x)| &\leqslant C((n^{-\frac{1}{2}}\frac{\delta_n(x)}{\phi(x)})^{\alpha_0} \\ +\min\{n^{\frac{r}{2}}\frac{\phi^r(x)}{\varphi^r(x)}, n^r\phi^r(x)\}h^r\omega_\phi^r(f,\delta)_{\bar{w}} &+ h^r\delta^{-r}\omega_\phi^r(f,\delta)_{\bar{w}}) \\ &\leqslant C((n^{-\frac{1}{2}}\frac{\delta_n(x)}{\phi(x)})^{\alpha_0} \\ + h^rM^r(n^{-\frac{1}{2}}\frac{\varphi(x)}{\phi(x)} \ + \ n^{-\frac{1}{2}}\frac{n^{-1/2}}{\phi(x)})^{-r}\omega_\phi^r(f,\delta)_{\bar{w}} &+ h^r\delta^{-r}\omega_\phi^r(f,\delta)_{\bar{w}}) \\ &\leqslant C((n^{-\frac{1}{2}}\frac{\delta_n(x)}{\phi(x)})^{\alpha_0} \\ +h^rM^r(n^{-\frac{1}{2}}\frac{\delta_n(x)}{\phi(x)})^{-r}\omega_\phi^r(f,\delta)_{\bar{w}} &+ h^r\delta^{-r}\omega_\phi^r(f,\delta)_{\bar{w}}). \end{aligned}$$

When $n \geqslant 2$, we have

$$n^{-\frac{1}{2}}\delta_n(x) < (n-1)^{-\frac{1}{2}}\delta_{n-1}(x) \leqslant \sqrt{2}n^{-\frac{1}{2}}\delta_n(x),$$

Choosing proper $x,\ \delta,\ n \in N$, so that

$$n^{-\frac{1}{2}}\frac{\delta_n(x)}{\phi(x)} \leqslant \delta < (n-1)^{-\frac{1}{2}}\frac{\delta_{n-1}(x)}{\phi(x)},$$

Therefore

$$|\bar{w}(x)\Delta_{h\phi}^r f(x)| \leqslant C\{\delta^{\alpha_0} + h^r\delta^{-r}\omega_\phi^r(f,\delta)_{\bar{w}}\}.$$

Which implies

$$\omega_\phi^r(f,t)_{\bar{w}} \leqslant C\{\delta^{\alpha_0} + h^r\delta^{-r}\omega_\phi^r(f,\delta)_{\bar{w}}\}.$$

So, by Berens-Lorentz lemma in [5], we get

$$\omega_\phi^r(f,t)_{\bar{w}} \leqslant Ct^{\alpha_0}.$$

□

REFERENCES

[1] P. L. Butzer, Linear combinations of Bernstein polynomials, Canad. J. Math. 5 (1953), pp. 559-567.

[2] H. Berens and G. Lorentz, Inverse theorems for Bernstein polynomials, Indiana Univ. Math. J. 21 (1972), pp. 693-708.

[3] Z. Ditzian, A global inverse theorem for combinations of Bernstein polynomials, J. Approx. Theory 26 (1979), pp. 277-292.

[4] D. Della Vechhia, G. Mastroianni and J. Szabados, Weighted approximation of functions with endpoint and inner singularities by Bernstein operators, Acta Math. Hungar. 103 (2004), pp. 19-41.

[5] Z. Ditzian and V. Totik, Moduli of Smoothness, Springer-Verlag, Berlin, New York (1987).

[6] M. Felten, Direct and inverse estimates for Bernstein polynomials, Constr. Approx., 14, 459-468.

[7] S. S. Guo, C. X. Li and X. W. Liu, Pointwise approximation for linear combinations of Bernstein operators, J. Approx. Theory 107 (2000), pp. 109-120.

[8] S. S. Guo, H. Tong and G. Zhang, Pointwise weighted approximation by Bernstein operators, Acta Math. Hungar. 101 (2003), pp. 293-311.

[9] G. G. Lorentz, Bernstein Polynomial, University of Toronto Press, Toronto (1953).

[10] W. M. Lu, L. Zhang and M.Y. Chai, Weighted approximation of functions with endpoint singularities by combinations of Bernstein operatorsarXiv: 1007.5436v2 [math.FA] 2 Aug 2010

[11] L. S. Xie, Pointwise simultaneous approximation by combinations of Bernstein operators, J. Approx. Theory 137 (2005), pp. 1-21.

[12] L. S. Xie, The saturation class for linear combinations of Bernstein operators, Arch. Math. 91 (2008), pp. 86-96.

[13] D. S. Yu, Weighted approximation of functions with singularities by combinations of Bernstein operators, J. Applied Mathematics and Computation. 206(2008),pp.906-918.

[14] D. S. Yu and D. J. Zhao, Approximation of functions with singularities by truncated Bernstein operators, Southeast Bull. Math. 30 (2006), pp. 1178-1189.

[15] D. X. Zhou, Rate of convergence for Bernstein operators with Jacobi weights, Acta Math. Sinica 35 (1992), pp. 331-338.

[16] D. X. Zhou, On smoothness characterized by Bernstein type operators, J. Approx. Theory 81 (1994), pp. 303-315.

[17] J. J. Zhang, Z. B. Xu, Direct and inverse approximation theorems with Jacobi weight for combinations and higer derivatives of Baskakov operators(in Chinese), Journal of systems science and mathematical sciences. 2008 28 (1), pp. 30-39.

Wen-ming Lu
School of Science, Hangzhou Dianzi University, Hangzhou, People's Republic of China
E-mail address: `lu_wenming@163.com`

Lin Zhang
Department of Mathematics, Zhejiang University, Hangzhou, People's Republic of China
E-mail address: `godyalin@163.com;linyz@zju.edu.cn`